

The Variable Outflow from the Chukchi Shelf to the Arctic Ocean

Thomas Weingartner
Institute of Marine Science
School of Fisheries and Ocean Sciences
University of Alaska
Fairbanks, AK 99775-7220
phone: 907-474-7993 fax: 907-474-7204 e-mail: weingart@ims.alaska.edu

Award Number: N00014-02-1-0308
<http://psc.apl.washington.edu/HLD/SBI2003/SBIHealy2003.html>
<http://www.whoi.edu/arcticedge.html>

LONG TERM GOALS

My long-term goal is to understand the mean and time-varying density structure and circulation dynamics of the Chukchi/Beaufort continental slope of the Arctic Ocean and how the slope interacts with the adjacent shelves and basin.

OBJECTIVES

The field effort and data analyses are addressing the following questions and objectives:

1. What are the characteristics of the shelf outflow from the Chukchi Sea to the Arctic Ocean?
2. What is the variability of these outflows and their water properties?
3. Which upstream processes are principally responsible for the variability?
4. contribute to the SBI goals by illuminating the mechanisms of shelf-basin exchange; and
5. promote further improvements in the rapidly growing array of models of arctic circulation, hydrographic structures, and variability by providing measurements against which to test the fidelity of these models.

The measurements include moored time series, supplemented by hydrographic and other ship-borne data. Our focus is both on the transformation and outflow from the shelf of Pacific waters that have entered through Bering Strait [*Weingartner et al.*, 2005; *Woodgate et al.*, 2005a; b; *Pickart et al.*, 2005], and on providing a context in which the SBI process studies can be embedded.

APPROACH

Our effort is part of a broader, interdisciplinary effort being undertaken by the NSF-ONR Shelf-Basin Interactions (SBI) project as part of NSF's global change program, ARCSS (Arctic System Science). My specific program is a co-ordinated effort with Knut Aagaard and Rebecca Woodgate (both at the University of Washington). We deployed 4 oceanographic moorings on the Chukchi Shelf and slope

| Report Documentation Page | | | | Form Approved OMB No. 0704-0188 | |
|--|------------------------------------|-------------------------------------|---|---|---------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE 30 SEP 2006 | | 2. REPORT TYPE | | 3. DATES COVERED 00-00-2006 to 00-00-2006 | |
| 4. TITLE AND SUBTITLE The Variable Outflow from the Chukchi Shelf to the Arctic Ocean | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Alaska, Institute of Marine Science, School of Fisheries and Ocean Sciences, Fairbanks, AK, 99775-7220 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT Same as Report (SAR) | 18. NUMBER OF PAGES 6 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

[Barrow Canyon (ca. 80 m depth), the Central Channel (ca. 50 m), and on the north-central shelf downstream of Herald Valley (ca. 70 m and 110 m)]. The moorings were initially deployed from the USCG icebreaker Polar Star in September 2002, re-covered and re-deployed in September 2003, with the final recovery cruise conducted in September-October 2004. The shelf moorings are examining the flow and thermohaline properties of the shelf and the slope moorings will measure these characteristics near the shelfbreak of the northern Chukchi Sea and in the western Beaufort Sea. Each mooring includes ADCPs for measuring vertical profiles of currents and temperature-conductivity recorders for measuring temperature and salinity. The mooring deployment/recovery cruises also involve extensive CTD, bottle sampling, and vessel-mounted-ADCP transects over the Chukchi shelf and along transects that cross the Chukchi-Beaufort shelfbreak and slope. In closely related efforts, R. Pickart, WHOI maintained a mesoscale array on the Beaufort slope near 152°W, and J. Swift, SIO provided CTD/rosette sections to support the mooring work. These sections also included ADCP measurements that give detailed spatial information about the velocity field. To provide a context for the process studies and the overall SBI effort, we are also examining other data sets that uniquely contribute to understanding the modification of the Pacific inflow, to its role and fate in the Arctic Ocean, and to the larger role of arctic shelf-basin interaction in the global ocean.

WORK COMPLETED

Final data processing has been completed and all moored data are being prepared for submittal to JOSS/EOL for archiving, as designated by the SBI program.

In our analysis work we have made a considerable effort to synthesize a variety of data to provide a broad perspective on arctic shelf-basin. Our analysis efforts under this grant have resulted in five published papers, along with numerous presentations at national and international meetings. Several of the published results new this year are described below.

RESULTS

The effects of changes in the processing of freshwater in the Arctic and sub-Arctic are of global importance, and the SBI region is an integral part of this problem set. For example, the inflow of freshwater through Bering Strait is equivalent to about three-fourths of the total runoff into the Arctic Ocean [Woodgate *et al.*, 2006].

The annual cycle of freezing and melting of sea ice is of major importance to conditions on the arctic shelves, including the SBI region, and the ensuing shelf conditions also affect the adjacent basins. Much of the ice production has been thought to occur in coastal polynyas, where extremely saline waters may form [Aagaard *et al.*, 1985; Weingartner *et al.*, 2005]. To explore this problem set, we have examined conditions in the St. Lawrence Island polynya on the northern Bering shelf (Figure 1), which feeds the Bering Strait throughflow and downstream Chukchi Sea. Using the records from fourteen year-long instrumented moorings deployed south of St. Lawrence Island, along with oceanographic drifters, we have assessed the circulation over the central Bering shelf and the role of polynyas in forming and disseminating saline waters over the shelf. We have paid particular attention to evaluating the Gawarkiewicz and Chapman [1995] model of eddy production within coastal polynyas. Principal results of our analysis include: 1) The northern central shelf near-surface waters exhibit westward flow carrying low-salinity waters from the Alaskan coast in fall and early winter, with consequences for water mass formation and biological production. 2) Within the St. Lawrence polynya, the freshening effect of winter advection is about half as large as the salting effect of surface

brine flux resulting from freezing. 3) Brine production over the Bering shelf occurs primarily offshore, rather than within the polynyas. 4) We find little evidence for the geostrophic flow adjustment predicted by recent polynya models. 5) In contrast to the theoretical prediction that dense water from the polynya is carried offshore by eddies, we find negligible cross-shelf eddy density fluxes within and surrounding the polynya and very low levels of eddy energy that decrease from fall to winter, even though dense water accumulated within the polynya and large cross-shore density gradients develop. 6) It is possible that dense polynya water was advected downstream of our array before appreciable eddy fluxes materialized.




Figure 1: Composite SAR and AVHRR image of the St. Lawrence Island polynya on 9 January 1999, adapted from Drucker et al. [2003]. The polynya appears strongly wind-driven, with the winds and waves forcing the frazil ice into the long linear streaks characteristic of a Langmuir circulation. Mooring locations are plotted. SAR image copyright 1999 by the Canadian Space Agency.

IMPACT/APPLICATIONS

Our work on the northern Chukchi shelf is collaborative with K. Aagaard and R. Woodgate, UW; R. Pickart, WHOI, who is concentrating on the adjacent Beaufort shelf and slope; and with J. Swift, SIO, who has made hydrographic measurements throughout the SBI region. Additionally, we have ongoing collaborations in the region with E. Carmack and F. McLaughlin, IOS and S. Smith, L. Llinas, and D. Hansell of U. Miami.

Major goals of the SBI initiative are to understand the physical processes responsible for water mass modification over the arctic shelves and slopes, and for exchanges with the interior ocean, as well as to understand the variability of this system. Our reported project addresses these goals directly. In particular, we are quantifying the large variability found in the Pacific-origin waters that flush the western Arctic shelves, as well as illuminating the origin of this variability. Much of the latter is generated in the Bering Sea [Woodgate *et al.*, 2005a], although the northward-flowing waters may in some years be further modified in the Chukchi, particularly during winter along the Alaskan coast [Weingartner *et al.*, 2005]. The shelf waters are subsequently discharged into the Arctic Ocean, where their seasonal and interannual variability are propagated long distances, in part by long-lived eddies that drift into the interior [Newton *et al.*, 1974; Manley and Hunkins, 1985], in part by topographically steered boundary currents that rim both the Polar Basin and its major ridge structures [Aagaard, 1989], and in part by other features of the circulation. This propagation leads to variability in regions far from the originating shelves [cf., Swift *et al.*, 1997 and Woodgate *et al.*, 2001 for examples]. An understanding of these effects and processes is vital to realistically modeling the Arctic Ocean and its global connections [Huang and Schmitt, 1993; Wadley and Bigg, 2002; DeBoer and Nof, 2004]].

Our measurements of the time-dependent shelf circulation provide important guidance to investigations of shelf productivity and biochemical cycling. For example, water parcels leaving the northern Chukchi shelf will exhibit substantially different carbon and nutrient loading, depending on their upstream trajectories and the season [Walsh *et al.*, 1997; Woodgate *et al.*, 2005a]. Our work also addresses other prominent issues, including the role of polynyas [Danielson *et al.*, in press]; the flux and processing of freshwater [Woodgate and Aagaard, 2005; Peterson *et al.*, 2006; Woodgate *et al.*, 2006; Aagaard *et al.*, in press; Serreze *et al.*, in press]; mixing over the slope [Woodgate *et al.*, 2005b]; and the needs of a variety of arctic simulations for accurate long-term boundary conditions and observations for verification.

IMPACT/APPLICATIONS

Our work is providing the observations and interpretations required to understand how Pacific waters influence the structure and dynamics of the Arctic Ocean. In addition we are providing other SBI investigators with ancillary information necessary for interpreting biological and geochemical data collected during the SBI program. Our measurements will also provide data sets essential for evaluating and testing the numerical model activities being undertaken in SBI.

TRANSITIONS

All SBI investigators will use our data and interpretations in examining their data sets.

RELATED PROJECTS

Our SBI project complements moored observations and CTD sampling being collected in this region by other NSF-supported SBI participants as well as Japanese efforts led by K. Shimada (JAMSTEC) and Canadian sampling led by E. Carmack (Institute of Ocean Sciences).

REFERENCES

- Aagaard, K., A synthesis of the Arctic Ocean circulation, *Rapp. P.-V. Reun. Cons. Int. Explor. Mer*, 188, 11-22, 1989.
- Aagaard, K., J.H. Swift, and E.C. Carmack, Thermohaline circulation in the arctic mediterranean seas, *J. Geophys. Res.*, 90, 4833-4846, 1985.
- Aagaard, K., T.J. Weingartner, S.L. Danielson, R.A. Woodgate, G.C. Johnson, and T.E. Whitledge, Some controls on flow and salinity in Bering Strait, *Geophys. Res. Lett.*, in press.
- Danielson, S., K. Aagaard, T. Weingartner, S. Martin, P. Winsor, G. Gawarkiewicz, and D. Quadfasel, The St. Lawrence polynya and the Bering shelf circulation: New observations that test the models, *J. Geophys. Res.*, 111, C09023, doi:10.1029/2005JC003268, 2006 [published, refereed]
- DeBoer, A. M., and D. Nof, The exhaust valve of the North Atlantic, *J. Clim.*, 17 (3), 417-422, 2004.
- Gawarkiewicz, G. and D. C. Chapman, A numerical study of dense water formation and transport on a shallow, sloping continental shelf, *J. Geophys. Res.*, 100, 4489-4508, 1995.
- Huang, R. X., and R. W. Schmitt, The Goldsbrough-Stommel circulation of the world ocean, *J. Phys. Oceanogr.*, 23, 1277-1284, 1993.
- Manley, T.O., and K. Hunkins (1985), Mesoscale eddies in the Arctic Ocean, *J. Geophys. Res.*, 90, 4911-4930.
- Newton, J.L., K. Aagaard, and L.K. Coachman (1974), Baroclinic eddies in the Arctic Ocean, *Deep Sea Res.*, 21, 707-719, 1974.
- Pickart, R.S., T. Weingartner, L.J. Pratt, S. Zimmermann, and D. J. Torres, Flow of winter-transformed Pacific water into the western Arctic. *Deep-Sea Res. II*, 52 (24-26), 3175-3198, 2005.
- Serreze, M.C., A.P. Barrett, A.G. Slater, R.A. Woodgate, K. Aagaard, R. Lammers, M. Steele, R. Moritz, M. Meredith, and C.M. Lee, The large-scale freshwater cycle of the Arctic, *J. Geophys. Res.*, in press.
- Swift, J.H., E.P. Jones, K. Aagaard, E.C. Carmack, M. Hingston, R.W. Macdonald, F.A. McLaughlin, and R.G. Perkin, Waters of the Makarov and Canada basins, *Deep-Sea Res. Part II*, 44, 1503-1529, 1997.
- Wadley, M. R., and G. R. Bigg , Impact of flow through the Canadian Archipelago and Bering Strait on the North Atlantic and Arctic circulation: An ocean modelling study, *Quart. J. Royal Met. Soc.*, 128, 2187-2203, 2002.

Walsh, J.J., D.A. Dieterle, F.E. Muller-Karger, K. Aagaard, A.T. Roach, T.E. Whitledge, and D. Stockwell, CO₂ cycling in the coastal ocean. II. Seasonal organic loading of the Arctic Ocean from source waters in the Bering Sea, *Continental Shelf Res.*, 17, 1-36, 1997.

Weingartner, T., K. Aagaard, R. Woodgate, S. Danielson, Y. Sasaki, and D. Cavalieri, Circulation on the north central Chukchi Sea shelf, *Deep-Sea Res. II*, 52 (24-26), 3150-3174, 2005.

Woodgate, R.A., and K. Aagaard, Revising the Bering Strait freshwater flux into the Arctic Ocean, *Geophys. Res. Lett.*, 32, L02602, doi:10.1029/2004GL021747, 2005.

Woodgate, R.A., K. Aagaard, and T. Weingartner, A year in the physical oceanography of the Chukchi Sea: Moored measurements from autumn 1990-1991, *Deep-Sea Res. II*, 52 (24-26), 3116-3149, 2005a.

Woodgate, R.A., K. Aagaard, and T.J. Weingartner, Interannual changes in the Bering Strait fluxes of volume, heat and freshwater between 1991 and 2004, *Geophys. Res. Lett.*, 33, L15609, doi:10.1029/2006GL026931, 2006.

Woodgate, R.A., K. Aagaard, J.H. Swift, K. Falkner, and W.M. Smethie Jr., Pacific ventilation of the Arctic Ocean's lower halocline by upwelling and diapycnal mixing over the continental margin, *Geophys. Res. Lett.*, 32, No. 18, L18609, doi:10.1029/2005GL023999, 2005b.

Woodgate, R.A., K. Aagaard, R.D. Muench, J. Gunn, G. Björk, B. Rudels, A.T. Roach, and U. Schauer, The Arctic Ocean boundary current along the Eurasian slope and the adjacent Lomonosov Ridge: Water mass properties, transports and transformations from moored instruments, *Deep-Sea Res. I*, 48, 1757-1792, 2001.

PUBLICATIONS

Aagaard, K., T.J. Weingartner, S.L. Danielson, R.A. Woodgate, G.C. Johnson, and T.E. Whitledge, Some controls on flow and salinity in Bering Strait, *Geophys. Res. Lett.* [in press, refereed]

Danielson, S., K. Aagaard, T. Weingartner, S. Martin, P. Winsor, G. Gawarkiewicz, and D. Quadfasel, The St. Lawrence polynya and the Bering shelf circulation: New observations that test the models, *J. Geophys. Res.*, 111, C09023, doi:10.1029/2005JC003268, 2006 [published, refereed]

Pickart, R.S., T. Weingartner, L.J. Pratt, S. Zimmermann, and D. J. Torres, Flow of winter-transformed Pacific water into the western Arctic. *Deep-Sea Res. II*, 52 (24-26), 3175-3198, 2005 [published, refereed].

Weingartner, T., K. Aagaard, R. Woodgate, S. Danielson, Y. Sasaki, and D. Cavalieri, Circulation on the north central Chukchi Sea shelf, *Deep-Sea Res. II*, 52 (24-26), 3150-3174, 2005 [published, refereed]

Woodgate, R.A., K. Aagaard, and T. Weingartner, A year in the physical oceanography of the Chukchi Sea: Moored measurements from autumn 1990-1991, *Deep-Sea Res. II*, 52 (24-26), 3116-3149, 2005 [published, refereed]